

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
BEFORE OPENING OR USING



APPROVED
LABEL



Infopest



Verified

INITIATOR[®]

SYSTEMIC PLANT INSECTICIDE AND FERTILISER

ACTIVE CONSTITUENT: 200 g/kg IMIDACLOPRID

GROUP **4A** INSECTICIDE

For use in the protection against insect damage and to provide enhanced growth in eucalyptus trees, roses, azaleas, lilypillies, palms and magnolias.

Guaranteed Minimum Fertiliser Analysis:

Nitrogen (N) (as slow release)	3.85%
Nitrogen (N) as Ammonium Nitrate	1.75%
Total Nitrogen (N):	5.6%
Phosphorous (P) (as calcium phosphate)	2.6%
Phosphorous (P) (Slow release)	2.8%
Total Phosphorous:	5.4%
Potassium (K) (Slow release):	4.2%
Magnesium (Mg) (Slow release):	1.4%

750 g Net

(300 x 2.5 g
Tablets)

MN 79701306

LBL 790000/0000

DIRECTIONS FOR USE

PLANT	PEST	RATE	CRITICAL COMMENTS
Initiator tablets will control or significantly reduce damage caused by insects (described below) that attack containerised, newly planted and established plants. Through insect protection and fertiliser effects the tablet has been shown to improve survival and growth rates of newly established seedlings, containerized trees and shrubs, and newly-planted or established ornamental plants.			
Roses	Aphids	1 tablet per m of plant height.	Transplanted stock: Place tablet(s) in planting hole, immediately beneath root ball, at time of planting Established in-ground or potted plants: Create a hole in the soil approximately 5 cm deep, as close to the absorptive roots as possible (at least 10 cm away from stem of plant). Place tablet in hole and ensure tablet is completely covered with soil after placement. In plants greater than 1 m in height, where multiple tablets are required, space out the tablets evenly around the plant.
Azaleas (in-ground and potted)	Azalea lace bug		
Lilypillies	Psyllids		
Potted palms	Mealybugs		
Magnolias	Scale		
Eucalypt trees	Various defoliating insect pests including Chrysomelid larvae and adults (e.g. <i>Chrysophtharta</i> spp., Paropsis spp., <i>Cadmus</i> spp.), scarab pests (e.g. <i>Heteronyx dimidiata</i> and <i>H. crinitus</i>) Suppression of damage caused by Autumn gum moth (<i>Mnesampela privata</i>) and Light brown apple moth (<i>Epiphyas postvittana</i>) Stem damaging scarab pests e.g. adult African Black Beetle (<i>Heteronyx arator</i>) Sap-sucking pests e.g. Psyllids and Gum-tree scale Root-feeding scarab pests e.g. <i>Heteronyx elongatus</i>	1 tablet per seedling tree	Transplanted seedling stock: Place tablet in planting hole, immediately beneath or adjacent to root ball, at time of planting

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

Initiator tablets are formulated to release active ingredient slowly. Release of active ingredient is dependent on presence of adequate soil moisture. For best results, irrigate the plant after treatment, and ensure that the plant receives regular irrigation for at least 2-3 months after treatment.

FORESTRY APPLICATIONS

Once the seedling has been placed into the planting hole where an Initiator tablet has been applied there is no requirement for any deviation from normal planting practice. This tablet formulation has been shown to significantly improve growth and establishment rates of eucalypt plantations in areas where insect pests are significant limitations to growth and establishment.

PRODUCTION AND LANDSCAPE ORNAMENTS

Initiator tablets can be used as directed on the plant species described above in areas such as production and retail nurseries, home gardens and residential complexes; business, office and shopping complexes; golf courses, airports, cemeteries, parks, playgrounds, athletic fields, and other landscaped recreational areas.

INSECTICIDE RESISTANCE WARNING: GROUP **4A** INSECTICIDE

For insecticide resistance management Initiator is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Initiator and other Group 4A insecticides may exist through normal genetic variation in its target population. The resistant individuals can eventually dominate the insect population if Initiator or other Group 4A insecticides are used repeatedly. The effectiveness of Initiator on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of this product to control resistant insects. Initiator tablets may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative.

PRECAUTIONS: Do not allow stock to graze treated trees or pasture adjacent to trees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT leave tablets exposed above ground as they are dangerous to wildlife, particularly birds.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Triple rinse container before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. If recycling, replace lid and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves. Wash hands after use.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre (phone 13 11 26).

MATERIAL SAFETY DATA SHEET: Additional information is listed in the Material Safety Data Sheet.

EXCLUSION OF LIABILITY: This product must be used strictly as directed, and in accordance with instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 60391/625g-1.25 kg/0209

Initiator[®] is a Registered Trademark of Bayer

**FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY PHONE 1800 033 111**

Bayer Environmental Science
Bayer CropScience Pty. Ltd.
A.B.N. 87 000 225 022
391-393 Tooronga Rd
Hawthorn East, Vic 3123
Telephone: 1800 804 479
www.bayeres.com.au



Bayer Environmental Science
A Business Operation of
Bayer CropScience

Please insert 100%
13 digit barcode here
9319784801750