

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Pride

Insecticide

ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP **4A** INSECTICIDE

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Distributed by:

Amgrow Pty Ltd (Commercial Products Division)

ABN 22 069 900 456

Unit B2A, 3-29 Birnie Ave,

Lidcombe NSW 2141

TEL: 1800 631 008

www.ecofertiliser.com.au

© Pride is a registered trademark of Ecofertiliser Pty Ltd.

APVMA Approval No. 64900/20L/0410



CONTENTS
20L

DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m ² Spray with at least 400 L water per hectare to ensure even coverage.	Apply at peak egg hatch, which is mid spring to mid summer depending on species.
	Larvae of Billbug	Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.	Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.
Roses	Aphids	25 mL/100 L	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids Azalea lace bug Bronze orange bug Harlequin bug Citrus mealybug Greenhouse thrips Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L	Spray buds and flowers as needed.
	Longtailed mealybug	50 mL/100 L + surfactant	Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100 L	Spray at first sign and then a week later.
	Soft scales	25 mL/100 L	Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid	25 mL/100 L	Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid (<i>Jamella australiae</i>)	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water	Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.



SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE	CRITICAL COMMENTS
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height	Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant	Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20 L capacity.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water	Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.
Roses	Aphids	3.5 mL/2 L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION



GENERAL INSTRUCTIONS

GROUP

4A

INSECTICIDE

Insecticide Resistance Warning

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid,

Whitefly and Melon Thrips: DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

Confined environments such as glasshouses

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

Note on Ornamentals

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

PROTECTION OF LIVESTOCK

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.



PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

® Trademark of Ecofertiliser

APVMA Approval No. 64900/20LL/0410

